2019

PHYSICS

(Theory)

Full Marks: 70

Pass Marks: 21

Time: Three hours

Attempt all Questions.

The figures in the right margin indicate full marks for the questions.

Question Nos. 1 to 10, are "Very Short Answer" type questions carrying 1 mark each.

1.	Define relative permittivity of a medium on the basis of Coulomb's law.	1
2.	A long straight thin-walled copper tube of radius R carries a current I along	g its
	length. What is the magnetic field along the axis of the tube? Give reason.	. 1
3.	At a very high frequency of a.c. capacitor behaves as a conductor. Why?	1
4.	How can wattless current be obtained in an a.c. circuit?	1
5.	On what factor does the velocity of electromagnetic wave in a medi	um
	depend?	1
6	Define angular dispersion	1

1.	What is threshold frequency?	1
8.	What will happen if electron revolving around the nucleus comes to rest?	1
9.	What is mass defect?	1
10.	Give the bandwidth of TV signal for transmission.	1
	Question Nos. 11 to 20 are 'Short Answer Type-II' questions carrying 2 maeach.	rks
11.	A glass rod held by hand can be charged by rubbing it with a silk cloth but	ar
	iron rod cannot be charged in a similar way. Explain this reason.	2
12.	If two point charges $8 \mu C$ and $2 \mu C$ are separated by a distance 1 m in	air
	At what point on the line joining the two charges is the electric field intense zero?	sity 2
13.	A 6 ohm non insulated wire is bent 180° in the middle and the two halves	are
	twisted together. What will be its new resistance?	2
14.	Give two points of difference between diamagnetic and ferromagne	tic
	substances.	2
15.	In the given circuit, calculate phase difference between the current and the sup	ply
	voltage. $_{4\mu F}$ $_{2H}$ $_{500\Omega}$	2

16. How are gamma rays produced? Give one example of its uses.

2

17. Draw a neat and label diagram of human eye.


-

2

18. If the Young's double slit experiment is performed in water instead of air, what would happen to the interference pattern obtained in the experiment? Explain.

2

19. In the given common emitter configuration, the current gain of the transistor is100. Find the output voltage of the amplifier.2



20. For faster action n-p-n transistor is used. Explain why?

2

Question Nos. 21 to 27 are 'Short Answer Type-I' questions carrying 3 marks each.

- 21. What is meant by equipotential surface? Show that electric field is always perpendicular to the equipotential surface.

 1+2=3
- Derive an expression for self inductance of an air cored solenoid of length *l*,cross-section area A and having number of turns N.
- Show that lateral displacement depends upon the angle of incidence and thickness of the glass slab.

- Obtain the expression for de-Broglie wavelength of an electron when it is accelerated by a potential difference V.
- 25. Mention three points of difference between alpha and beta particles.
- 26. Explain with neat labelled circuit diagram, how zener diode is used as a voltage regulator?
- 27. Define the terms transducer, attenuation and repeater. $1 \times 3 = 3$

Question Nos. 28 to 30 are 'Long Answer Type' questions carrying 5 marks each.

28. What are e.m.f. and terminal potential difference of a cell? Derive the expressions for equivalent e.m.f. and internal resistance when two different cells are connected in parallel.

2+3=5

OR

State Kirchhoff's laws of an electrical network. Using the Kirchhoff's laws, deduce the condition for which Wheatstone Bridge is balanced. 2+3=5

29. Draw a schematic labelled diagram of a cyclotron. Deduce an expression for the cyclotron frequency and show that it does not depend on the speed of the charged particle.
1+3+1=5 Draw a neat labelled diagram of a moving coil galvanometer. Prove that the deflection produced in the galvanometer is directly proportional to the current flowing through it and define current sensitivity. 1+3+1=5

 Derive thin lens formula for a convex lens when real image is formed by using the necessary ray diagram.

A convex lens of refractive index 3/2 has a focal length of 20 cm in air. Calculate the change in its focal length when it is immersed in water of refractive index 4/3.

OR

Verify Snell's law of refraction using. Huygen's principle with the help of diagram.

The intensity of light for maxima and minima in an interference pattern is 100:64. Calculate the ratio of intensities of the coherent sources producing this pattern.

3+2=5

Question Nos. 31 to 34 are 'Multiple Choice Type' questions carrying 1 mark each. Choose the correct answer out of the four alternatives and rewrite the correct answer.

31. A coil having an area A is placed perpendicularly in a magnetic field which changes from B to 3B in time interval t, the emf induced in the coil is 1

- A. 0
- B. BA
- C. $\frac{BA}{t}$
- $D. \qquad \frac{2BA}{t}$
- 32. For constructive interference, path difference between the waves at the point of observation will be
 - A. odd multiple of $\frac{\lambda}{2}$
 - B. even multiple of $\frac{\lambda}{2}$
 - C. odd multiple of λ
 - D. even multiple of λ .
- 33. In which region of electromagnetic spectrum does the Balmer series of hydrogen atom belongs to?
 - A. visible
 - B. infra-red
 - C. ultraviolet
 - D. X-rays.

- A. loss of signal
- B. line of signal
- C. loss of sight
- D. line of sight.